### Cognition for Communication In PD: The Role of Speech Language Pathology

ZOE KRIEGEL, PH.D. CCC-SLP ASSISTANT PROFESSOR, DIVISION OF COMMUNICATION DISORDERS UNIVERSITY OF WYOMING

02 April, 2025

## **Today's Session**

**APPROACHING FEAR OF COGNITIVE CHANGES** 

**SPEECH LANGUAGE PATHOLOGY.. AND COGNITION?** 

2

COGNITIVE COMMUNICATION & PD

WHO IS INVOLVED?

5 **COGNITIVE-**COMMUNICATION **DIFFICULTIES AND STRATEGIES** 

6 LOOKING AHEAD WITH ACTION

Aging and living with Parkinson disease (PD) may bring about **fears of the future** and about the possibility of cognitive difficulties and dementia (Corner & Bond, 2004; Mantri et al., 2024)

 At least 10% of individuals living with PD have an express concern about developing dementia or cognitive impairment in the future (Mantri et al., 2024)

"I fear the future and am terrified of getting dementia"


- Stigma and fear of cognitive changes or dementia may act as a barrier for seeking help about these concerns (Parker et al., 2020)
- Facilitators toward seeking help about cognitive difficulties include:
  - Recognition of symptoms
  - Prior knowledge and contacts
  - Support from informal network

- Stigma and fear of cognitive changes or dementia may act as a barrier for seeking help about these concerns (Parker et al., 2020)
- Facilitators toward seeking help about cognitive difficulties include:
  - <u>Recognition of symptoms</u>
  - Prior knowledge and contacts
  - Support from informal network

- Stigma and fear of cognitive changes or dementia may act as a barrier for seeking help about these concerns (Parker et al., 2020)
- Facilitators toward seeking help about cognitive difficulties include:
  - Recognition of symptoms
  - Prior knowledge and contacts
  - Support from informal network

- Stigma and fear of cognitive changes or dementia may act as a barrier for seeking help about these concerns (Parker et al., 2020)
- Facilitators toward seeking help about cognitive difficulties include:
  - Recognition of symptoms
  - Prior knowledge and contacts
  - <u>Support from informal network</u>


## **TEAM FOR COGNITION**





9

## **TEAM FOR COGNITION**





## LANGUAGE & COGNITION

## Therapy is effective with a focus on: supporting access to life

## LANGUAGE & COGNITION

How are you going to clean up this room?

Think about the exact steps you would need to take in your home to get this room back in working order.



12

Difficulties with the ability to attend to or organize information and/or solve problems using thinking and language functions. These may affect:

- Selective Attention
- Word Finding (Language)
- Executive Functioning

## SELECTIVE ATTENTION

Difficulties maintaining focus on the desired content and ignoring unimportant information

#### **REDUCING COGNITIVE BURDEN**

- Select communication environments with fewer competing demands
- 1 conversation in a close space
- Remove radio, tv, background noise

#### LOOK HERE

## **WORD FINDING**

Difficulties retrieving the intended lexical item (word) when faced with a naming task or during conversation

- Exacerbated by acute stress
- What is your preference and when?

#### CIRCUMLOCUTION

- Using descriptive language or
- paraphrasing an idea when the
- precise word is difficult to recall.
- "Talking around the word" 15

## **WORD FINDING**

Difficulties retrieving the intended lexical item (word) when faced with a naming task or during conversation

- Exacerbated by acute stress
- What is your preference and when?

# JUSTICE

#### CIRCUMLOCUTION

- Using descriptive language or
- paraphrasing an idea when the
- precise word is difficult to recall.
- "Talking around the word" 16

## **EXECUTIVE FUNCTION**

A group of higher-level cognitive functions including planning, judgement, inhibition, and others

- Helps with decision-making and behavior
- May show up as difficulty planning steps and making the next decision for completing a task

#### **PURPOSEFUL PLANNING**

- Allowing a moment to identify the goal and the steps you intend to take to get there 17
- Task outline (written or verbal)

## LOOKING AHEAD & ACTION

- Baseline testing
  - The best way to detect subtle changes is to have a <u>clear starting point</u>
  - Ask your provider if you might have access to comprehensive neuropsychological testing
- Ensure that you are <u>talking with your provider</u> about potential fears, difficulties, and/or family history
- Ask your provider about SLP or OT therapy for <u>current</u> cognitive challenges you may be experiencing

## THANK YOU

Zoe Kriegel, Ph.D. CCC-SLP Assistant Professor, Division of Communication Disorders University of Wyoming Email: zkriegel@uwyo.edu Ph: 307-766-6090

## REFERENCES

Corner, L., & Bond, J. (2004). Being at risk of dementia: Fears and anxieties of older adults. Journal of aging studies, 18(2), 143-155.

Mantri, S., Purks, J. L., Kinel, D., Arbatti, L., Hosamath, A., Allen, A., ... & Marras, C. (2024). In Their Own Words: Fears Expressed by People with Parkinson's Disease in an Online Symptom Database. Journal of Parkinson's Disease, 14(4), 865-872.

Parker, M., Barlow, S., Hoe, J., & Aitken, L. (2020). Persistent barriers and facilitators to seeking help for a dementia diagnosis: a systematic review of 30 years of the perspectives of carers and people with dementia. International psychogeriatrics, 32(5), 611-634.

20